




Marine Recreational Information Program

Transition to Improved Survey Designs

Dave Van Voorhees Kelly Denit

Estimating Total Recreational Catch





Improved Survey Designs

Fishing Effort Survey

- New mail survey of shore & private boat trips
- Replaces legacy telephone survey
- Uses USPS database and angler registries
- Higher, more accurate estimates of trips

Access Point Angler Intercept Survey

- New design for sampling catch
- Better time-of-day coverage
- More statistically sound



Shore

Private Boats

Fishing Effort Survey









Improved questionnaire.



More complete answers.



Extensive Testing and Peer Review

"The [Fishing Effort Survey] methodologies, including the address-based sampling survey design, are major improvements."

"The current methods used in the APAIS ... are a vast improvement over the previous sampling and estimation procedures and reflect state of the art methods in survey sampling."

> The National Academies of




SCIENCES

Transition Plan

Improving designs can change estimates Transition Plan developed by NOAA, states, councils, and commissions.



- Benchmarking (mail vs. phone)
- Calibration (old vs. new)
- Convert "old" estimates into "new" currency
 - Stock assessments Management decisions



2018 Annual Catch Limits




2018 ACLs set using CHTS estimates

FES \leftarrow CHTS Calibration model converts both directions






Private Boat Effort





Shore Effort





Private Boat Effort – Change Ratios

	Overall	1981-1999	2000-2014	2015-2017
Atlantic Coast	2.03	1.78	2.15	2.85
New England	1.83	1.65	1.92	2.34
Mid-Atlantic	1.98	1.71	2.13	2.91
South Atlantic	2.18	1.92	2.29	3.00




Shore Effort – Change Ratios

	Overall	1981-1999	2000-2014	2015-2017
Atlantic Coast	4.55	4.34	4.57	5.75
New England	3.91	3.72	4.00	4.61
Mid-Atlantic	3.80	3.37	3.98	5.33
South Atlantic	5.24	5.13	5.18	6.30




Bluefish Harvest



NOAA FISHERIES

U.S. Department of Commerce | National Oceanic and Atmospheric Administration | NOAA Fisheries | Page 14


Bluefish Total Catch



NOAA FISHERIES


U.S. Department of Commerce | National Oceanic and Atmospheric Administration | NOAA Fisheries | Page 15

Red Drum Harvest







Red Drum Total Catch





















Black Sea Bass Total Catch

























Atlantic Cod Harvest



NOAA FISHERIES


U.S. Department of Commerce | National Oceanic and Atmospheric Administration | NOAA Fisheries | Page 24

Atlantic Cod Harvest






Atlantic Cod Total Catch



NOAA FISHERIES

U.S. Department of Commerce | National Oceanic and Atmospheric Administration | NOAA Fisheries | Page 26

Atlantic Cod Total Catch



NOAA FISHERIES

U.S. Department of Commerce | National Oceanic and Atmospheric Administration | NOAA Fisheries | Page 27

Impacts on Recreational Fishing







Stock Assessment Schedule

Species	Region	Туре	Timing
Striped Bass	Atlantic Coast	Benchmark	Fall 2018
Summer Flounder	Mid-Atlantic Coast	Benchmark	Fall 2018
Red Snapper	Gulf of Mexico	Data Update	Fall 2018
Atlantic Cod	Gulf of Maine	Operational	Early 2019
Black Sea Bass	Mid-Atlantic Coast	Operational	Early 2019
Bluefish	Atlantic Coast	Operational	Early 2019
Scup	Atlantic Coast	Operational	Early 2019
Spot	Atlantic Coast	Operational	Early 2019
Red Porgy	Southern Atlantic Coast	Full Update	May 2019
Greater Amberjack	Southern Atlantic Coast	Full Update	May 2019
Red Grouper	Gulf of Mexico	Full Update	July 2019
Gray Triggerfish	Gulf of Mexico	Full Update	Aug. 2019
Yellowtail Snapper	Southern Atlantic Coast/Gulf of Mexico	Benchmark	Nov. 2019
Red Snapper	South Atlantic	Full Update	2020



Key Takeaways

- We see a substantial increase in effort, especially in the shore mode.
- Those stocks that have high proportion of catch from shore are more heavily impacted.
- The changes in effort are generally larger in more recent years, mostly driven by the "wireless" effect.
- 2018 catch will be back-calibrated to ensure ACLs and catch are in the same "currency."
- Stock assessments incorporating this new data will be used to determine stock status and ACLs.







What's Next

2018

- Revised total catch estimates now available for use in planned stock assessments.
- 2018 ACLs and catch estimates will be in the same "currency."
- Preliminary management changes may be made for reassessed stocks.
 - Calibrated statistics incorporated into additional stock assessments.

2020

2019

• Based on new stock assessments, management changes could occur for a number of species.



Upcoming Presentations



August 14

Mid-Atlantic Fishery Management Council



September 24

- New York State Department of Environmental Conservation,
 - Division of Marine Resources, Marine Resources Advisory Council



Scheduling Underway



September 5-6

NOAA Fisheries Atlantic Highly Migratory Species Advisory Panel



September 20

• South Atlantic Fishery Management Council



September 24

• New York State Department of Environmental Conservation, Division of Marine Resources, Marine Resources Advisory Council



September 25-27

Fishery Management Council New England Fishery Management Council



Scheduling Underway



October 2-4

 Gulf of Mexico Fishery Management Council Scientific & Statistical Committee



October 15-18

Gulf States Marine Fisheries Commission



October 22-25

• Gulf of Mexico Fishery Management Council







Questions?

