

ASMFC Alternative Management Plan for Shad and River Herring in Florida

Prepared by

Reid Hyle

Approved by the Shad and River Herring Management Board

August 6, 2025

On Behalf of

Florida Fish and Wildlife Conservation Commission

Fish and Wildlife Research Institute and Division of Marine Fisheries Management

This alternative management plan for American Shad and Blueback Herring in Florida addresses American Shad management outside of the St. Johns River System and Blueback Herring in all state waters.

System Descriptions

The St. Johns River in Florida drains 22,900 km² along east central Florida from Vero Beach to Jacksonville. The primary spawning runs of American Shad (*Alosa sapidissima*) and Blueback Herring (*Alosa aestivalis*) in Florida historically were and currently are in the St. Johns River. Spawning of *Alosa* spp. occurs from late December to early May in most years, with peak activity from mid-January to mid-March for American Shad and Blueback Herring (Walburg 1960, Williams and Bruger 1972, Williams et al. 1975, McBride and Holder 2008, McBride et al. 2010). The spawning grounds of American Shad have been documented from rkm 230 to rkm 433 near the headwaters (Williams and Bruger 1972, Williams et al. 1975). Of that distance 160km can be classified as river and 43 km as lake. Primary spawning grounds of American Shad were in river habitats between rkm 275 and rkm 360 (Williams and Bruger 1972). Contemporary egg collection (Miller et al. 2012b) and telemetry (Dutterer et al. 2011) confirm that American Shad spawning grounds still exist between rkm 230 and a weir at rkm 415. Blueback Herring spawning area overlap American Shad and may extend further downstream, but the specific habitats have not been identified (Williams et al. 1975). The mainstem run of the St. Johns River supported significant commercial fisheries of shad and river herring in the 19th and 20th centuries and continues to support a small recreational fishery for American Shad but not Blueback Herring.

Other Atlantic Coast Systems North of Cape Canaveral

The Nassau River is a small river restricted to the coastal plain between the St. Marys River and the St. Johns River. It has a drainage area of ~1,000 square miles (ACOE 1999). There is a passing reference to “a few fish” being taken from the Nassau River in Walburg and Nichols 1967 and no contemporary records of shads being taken in the Nassau River. Most of the stream is under tidal influence. There are no contemporary records of *Alosa* spp. in the Nassau River.

Pelicer Creek and the Tomoka River are small coastal streams with drainages areas of 412 and 385 km² respectively and stream lengths of <16 km. They are considered part of the “Northern Coastal Basin” that drain into a shared lagoon (SJRWMD 2003, Brown and Orel 1995). Neither received mention of having *Alosa* fisheries in the mid-20th century federal studies e.g. Walburg and Nichols 1967 and Williams and Grey 1975. Rulifson et al. 1982 extended the probable range of *Alosa* as far south as the Tomoka River. That finding was based on asking then Florida Game and Freshwater Fish Commission biologists and verbal records mentioned in Williams and Grey 1975. No specimens were recorded or vouchered and no quantity of fish or confirmation of spawning of *Alosa* in these small systems south of the St. Johns River have been documented. A faunal survey in the 1990s that recorded 59 species of fish in the Tomoka River did not record any *Alosa* species.

American Shad and Blueback Herring appear to be functionally absent from the Nassau River, Pelicer Creek, and the Tomoka river. Pelicer Creek and Tomoka River are likely outside the natural range of American Shad and Blueback Herring.

Florida Blueback Herring Fisheries

There has not been a fishery for Blueback Herring (*Alosa aestivalis*) in Florida for more than 30 years. Blueback Herring were likely an important commercial fishery in Florida in the 19th and early 20th centuries but catch data from that time period are unreliable. Landings of ‘alewife’ were reported up to a peak of around 1 million pounds in the early 20th Century. However, ‘alewife’ were often the combined landings of Blueback Herring, Hickory Shad (*Alosa mediocris*), and Menhaden (*Brevoortia* spp.). It is unclear what proportion of the landings was herring, though herring were harvested and salted for market at the time. By the mid-20th Century, herring harvest was limited to bycatch in other fisheries and was sold as crab and catfish bait (Williams et al., 1975). Those bycatch fisheries eventually ended following various gear restrictions in Florida. Blueback Herring in Florida are not harvested by either commercial or recreational anglers and no harvest has been recorded since the 1960s. Almost all landings that did occur historically were in the St. Johns River. If a Blueback Herring Fishery were to develop it would be restricted to hook and line gear by existing regulations.

Florida American Shad Fisheries

American Shad (*Alosa sapidissima*) fishery and management history is described in the American Shad Sustainable Fishing Plan for the St. Johns River System, Florida. Commercial and recreational fishing for American Shad in Florida almost exclusively occurred in the St. Johns River for several decades. There has not been any commercial landing of American Shad reported from anywhere in Florida since 2000.

Recreational Fisheries American Shad Outside of the St. Johns River

No directed recreational fishing for American Shad or incidental catch of American Shad has been documented in Florida waters other than in the middle/upper St. Johns River and adjacent Econlockhatchee River (Walburg and Nichols 1967, Walburg, 1960, and Williams and Bruger 1972). MRIP surveys have not recorded American Shad fishing or catch to occur in the coastal systems (Nassau, Pelicer, Tomoka). FWC has regular contact with recreational fishing clubs (e.g. First Coast Fly Fishing Club, Mosquito Lagoon Fly Fishing Club, Orlando Kayak Fishing Community) from Jacksonville to Orlando, and none report fishing for or catching American Shad outside the St. Johns River system.

Regulations

There are no bag or size limits for Blueback Herring, but take is restricted to hook and line gear.

As of January 1, 1997 hook-and-line gear was the only permissible gear for shad (American, Hickory, and Alabama) and river herring (blueback herring) in Florida (46-52.001(2), FAC and later updated to 68B-52.006, FAC). There was also a 10-fish aggregate bag limit implemented for shad (American, Hickory, and Alabama) at that time (46-52.001(3), FAC and later updated to 68B-52.004, FAC).

Fishery Dependent Monitoring

A creel survey is conducted annually on the St. Johns River. The survey is focused on American Shad but occurs in an area that overlaps Blueback Herring spawning grounds. There is also a creel survey from January through April that rotates between Lakes George, Monroe, and Crescent. These lakes are natural wide spots in the lower St. Johns River. No river herring catch, harvest, or directed effort has been recorded in angler surveys outside the American Shad fishery on the Middle and Upper St. Johns River. There is one recent instance of Blueback Herring being reported in a creel survey from a few anglers fly fishing for American Shad in 2018. However, these anglers reported them as 'baby shad' and promptly released their catch. The year of this reported sighting of blueback herring coincidentally coincided with a year of high abundance recorded in the electrofishing surveys (Figure 1). There is no Fishery Dependent Monitoring by FWC on the St. Marys River or the other coastal systems; Nassau River, Pellicer Creek, and Tomoka River. MRIP surveys have not recorded any Blueback Herring fishing or catch in the small coastal systems, St. Marys, or St. Johns River. The FWC creels in the St. Johns and MRIP should continue indefinitely.

Fishery Independent Monitoring

FWC conducts spawning stock and juvenile sampling for American Shad in the St. Johns River. Both of these surveys encounter Blueback Herring. These surveys could produce a CPUE based abundance index for both life stages. The spawning stock survey is a standardized electrofishing survey from January through March and the time series is 2003 through the present (Figure 1). The juvenile survey is a standard pushed trawl that runs bi-weekly from March through July. These data are reported in the annual compliance report to ASMFC (Figure 2). The pushnet is effective at capturing YOY Blueback Herring (Figure 2). Georgia DNR conducts a standardized electrofishing survey on the St. Marys River that has not to date encountered Blueback Herring. There are no credible records of Blueback Herring runs in the Nassau River, Pellicer Creek, or Tomoka River so directed fishery independent sampling there is not planned.

Management Alternative

Florida seeks to leave the current regulations for shad and river herring in the Florida Administrative Code unchanged until either 1) there is evidence that harvest of Blueback Herring is occurring anywhere in the state; or 2) there is evidence that harvest of American Shad outside the monitored St. Johns River system is occurring.

No commercial gears that could result in incidental catch of shad or river herring, such as pound nets, gill nets, or haul seines, are operating in Florida waters. No recreational fisheries are known to be

catching or harvesting Blueback Herring anywhere in Florida or American Shad outside of the monitored St. Johns River.

Blueback Herring and American Shad are absent or functionally absent from all drainages in Florida except the St. Johns River system. Therefore, it is not possible to develop useful metric of sustainability in systems outside the St. Johns. Monitoring in the St. Johns River could yield index-based benchmarks for Blueback Herring similar to those for American Shad if needed. However, the absence of any harvest, directed catch, or significant incidental catch preclude there being any conservation benefit to changing regulations to prohibit the harvest of Blueback Herring or American Shad beyond the existing gear and bag restrictions.

FWC proposes to continue its existing fishery dependent and fishery independent monitoring that focus on the St. Johns River where there are known populations of American Shad and Blueback Herring and where there is a monitored recreational fishery for American Shad and any incidental capture of other *Alosa* species. FWC will rely on the American Shad creel survey and other angler creel surveys to monitor for the existence of Blueback Herring catch or harvest in the St. Johns River. FWC will rely on MRIP and contact with recreational fishing organizations to detect *Alosa spp.* recreational catch in waters outside of the St. Johns River Basin. FWC will keep informed of GADNR monitoring of the St. Marys River for information about Blueback Herring or American Shad harvest and coordinate a response with Georgia DNR if data warrant. FWC will add reporting of data collected in accordance with this alternative management plan to its annual Shad and River Herring FMP compliance report.

Triggered Action

If any source detects non-zero Blueback Herring harvest anywhere in Florida for three consecutive years or there is recorded American Shad harvest outside the St. Johns River basin for three consecutive years, then Florida FWC will initiate a process to demonstrate sustainability for the affected population through monitoring compliant with Amendment 3 and begin the process to implement appropriate management actions. Since the American Shad bag limit already applies throughout Florida, one potential action could be to incorporating Blueback Herring into the 10 fish aggregate bag limit if there are indications of harvest for three consecutive years.

Literature Cited

- ACOE 1999. Nassau River Basin Comprehensive Floodplain Management Study. Special Publication SJ99-SP7. U.S. Army Corps of Engineers.
- Brown, M.T., and J. Orell. 1995. Tomoka River and Spruce Creek Riparian Habitat Protection Zone. Final Report. St. Johns River Water Management District Contract 94K353.

- Dutterer, A.C., Allen, M.S., and W.E. Pine. 2011. Spawning Habitats for American Shad as the St. Johns River, Florida: Potential for Use in Establishing MFLs. Special Publication SJ2012-SP1. St. Johns River Water Management District. Palatka, FL.
- McBride, R.S., Harris, J.E., Hyle, A.R., and J.C. Holder. The Spawning Run of Blueback Herring in the St. Johns River, Florida. Transactions of the American Fisheries Society 139(2): 598-609.
- McBride, R.S and J. C. Holder. 2008. A review and updated assessment of Florida's anadromous shads: American shad and hickory shad. North American Journal of Fisheries Management 28: 1668-1686.
- McLEAN, W. M. 1955. The fishes of the St. Johns River system. Ph.D. dissertation. University of Florida, Gainesville. 362 pp.
- Miller, S.J., Brockmeyer, R.E., Tweedale, W., Shenker, J., Keenan, L.W., Connors, S., Lowe, E.F., Miller, J., Jacoby, C., and L. McCloud. 2012b. Appendix 12.C. Potential Withdrawal Effects on Anadromous Herrings. St. Johns River Water Supply Impact Study. St. Johns River Water Management District. Technical Publication SJ2012-1.
- RULIFSON, R. A., M. T. HUIH, AND R. W. THOESSEN. 1982a. Anadromous fish in the southeastern United States and recommendations for development of a management plan. United States Fish and Wildlife Services, Fishery Resource Region 4. Atlanta
- SJRWMD 2003. Northern Coastal Basin Surface Water Improvement and Management Plan. St. Johns River Water Management District. Palatka, FL.
- Walburg, C. H., and P. R. Nichols. 1967. Biology and management of the American shad and status of the fisheries, Atlantic coast of the United States, 1960. U.S. Fish and Wildlife Special Scientific Report Fisheries. 550:1-105
- Walburg, C. H. 1960. Abundance and life history of shad, St. Johns River, Florida. Fishery Bulletin, U.S. 60: 486-501.
- Williams, R.O., and G.E. Bruger. 1972. Investigations on American shad in the St. Johns River. Florida Department of Natural Resources Marine Research Laboratory Technical Series 66. 1-49
- Williams, R. O., W. F. Grey, and J. A. Huff. 1975. Study of anadromous fishes of Florida. Completion Report for the period 1 May 1971 to 30 June 1974 for research funded by the Anadromous Fish Act (PL 89-304). National Marine Fisheries Service, St. Petersburg, Florida




Figure 1. Annual geometric mean electrofishing catch per transect of Blueback Herring from the St. Johns River, Florida *Alosa* spawning stock survey. Each transect consisted of 10 minutes of electrofishing effort within a randomly selected 1km portion of the river.




Figure 2. Geometric mean Blueback Herring catch per unit effort from the annual Alosa pushnet survey in the lower St. Johns River, Florida.